

**CNPS Inventory of Rare and Endangered Plants of
California**

rareplants.cnps.org



PLANT DETAIL



Classification

Scientific Name	<i>Elodium blandowii</i> (F. Weber & D. Mohr) Eckel
Common Name	Blandow's bog moss
Family	Helodiaceae
Element Code Name	NBMUS3C011
USDA Plants Symbol	HEBL2
Synonyms/Other Names	<i>Helodium blandowii</i> (Web. & Mohr) Warnst.

Conservation Status

California Rare Plant Rank	2B.2
Global Rank	G4
State Rank	S2
CESA	None
FESA	None
Other Status	USFS_S
CRPR Changes	changed from 2B.3 to 2B.2 on 2021-05-26
Date Added	1/1/2001
Last Change	7/14/2021

Ecology and Life History

Lifeform	moss
Blooming Period	
Elevation: m (ft)	1862-2700 (6110-8860)
General Habitat	Meadows and seeps, Subalpine coniferous forest
General MicroHabitat	Damp soil
Micro Habitat	

Element Occurrence Data from California Natural Diversity Database

Total Element Occurrences 16

Element Occurrence Ranks

Excellent (A)	2
Good (B)	3
Fair (C)	0
Poor (D)	0
None (X)	0
Unknown (U)	11

Occurrence Status

Historical, > 20 years	2
Recent, < 20 years	14

Presence

Presumed Extant	16
Possibly Extirpated	0
Presumed Extirpated	0

Location

CA Endemic No

Counties

El Dorado (ELD), Fresno (FRE), Mono (MNO), Siskiyou (SIS), Tulare (TUL), Tuolumne (TUO)

States

California (CA), Idaho (ID), Indiana (IN), Michigan (MI), Montana (MT), New York (NY), Oregon (OR), Vermont (VT), Washington (WA), Wisconsin (WI), Wyoming (WY)

Quads

Convict Lake (3711857), Freel Peak (3811978), Glass Mountain (3711876), Mt. Morgan (3711846), Mt. Patterson (3811943), North Palisade (3711815), Panther Rock (4112272), Sirretta Peak (3511883), Sonora Pass (3811936), South Lake Tahoe (3811988), Tennant (4112158)

Notes

Definitions of codes preceding a county and/or quad:

* Presumed extirpated

(*) Possibly extirpated

Species may be present in other areas where conditions are favorable. These data should NOT be substituted for pre-project review or for on-site surveys.

Notes

Threatened by grazing and trampling. Possibly threatened by hydrological alterations. Very distinctive due to abundant paraphyllia (minute leaf-like structures borne on the stems or branches among the leaves). See *Bot. Taschenb.* p. 332 (1807) for original description, and *Krypt. Fl. Brandenburg* 2:692 (1905) for revised nomenclature.

Threats

Taxonomy

Other

References