

Rare Plant Status Review: *Antennaria lanata*

Misa Ward (CNPS) and Roxanne Bittman (CNDDDB)

June 10, 2004

Antennaria lanata was reported to occur in California following the publication of the Jepson Manual. Although this taxon occurs in other western states and Canada, it is known in California from only a single locality. Ferlatte collected a specimen from the Trinity Alps (Siligo Meadows) in 1975. This specimen was annotated as *Antennaria lanata* by Dean Taylor and confirmed by Randall Bayer, the current author for the next version of the Jepson Manual

Please review the draft CNPS *Inventory* record below, and respond Yes or No on the proposal to add this taxon to the *Inventory* and CNDDDB. Please provide any input on the highlighted items in the record for which more information is needed. If responding No, please provide supporting information.

Recommended Actions

CNPS: Add to CNPS List 2

CNDDDB: Add to CNDDDB as G5 / S1.1

Draft CNPS *Inventory* record:

Antennaria lanata (Hook.)Greene

“woolly pussy-toes”

Asteraceae

List 2 / RED 3-3-1

Trinity, Oregon, Washington, and elsewhere

667B

Meadows and seeps / rocky; elevation 2225 meters.

Perennial herb, blooms July-August

Known in CA from only one occurrence at Siligo Meadows. Not in *The Jepson Manual*. See *Pittonia* 3:288 (1898) for revised nomenclature.

*[Habitat types are based on occurrences in other states...is this accurate for CA? The August blooming period assumes the CA specimen was flowering.]

Sent to NW